

Hip surveillance and management of the displaced hip in cerebral palsy

J. E. Robb · G. Hägglund

Received: 19 February 2013 / Accepted: 17 June 2013 / Published online: 18 August 2013
© EPOS 2013

Abstract

Introduction This article provides an overview of the management of a displaced hip in children with cerebral palsy and considers surveillance programmes, principles of surgical reconstruction and options for the salvage of an unreconstructable hip in these children.

Conclusion Hip dislocation in CP is potentially preventable if children are included from an early age in a surveillance programme that includes repeat radiographic and clinical examinations, and preventive treatment for hips that are displacing. A surveillance programme should be based on the child's age, GMFCS level and migration percentage (MP), and surgical prevention may be considered in children with a MP exceeding 33 %.

Keywords Cerebral palsy · Child · Hip displacement · Surveillance programme · Operative management

Introduction

It has been known for more than 50 years that hip dislocation in a child with cerebral palsy (CP) may be preventable [1]. Few countries have established screening programmes for the hip in patients with CP, although in many centres an annual pelvic radiograph is performed in

patients at risk of hip displacement. In practice, children often first present to a clinician with established radiological signs of hip displacement.

The cause of hip displacement in CP is not completely clear, but the hip is usually normal at birth. High hip joint reaction forces and abnormal orientations of force vectors are relevant and place an abnormal force on the acetabulum, which may then begin to deform [2]. The role of spasticity remains unclear, but is probably overestimated, because hip dislocations occur in hypotonic as well as hypertonic conditions. Another factor may be positioning [3], and many children with severe CP rest in the same position—particularly at night—in side-lying, where one hip remains in adduction. A further influence may be impaired proprioception, which results in inadequate muscle control [4].

A saucer-shaped deformity develops in the acetabulum, usually in a craniolateral direction [5], and the femoral head gradually migrates proximally and may also then become deformed. As the femoral head migrates proximally, it may be subjected to increasing forces on the medial side which result in medial flattening of the head. The lateral side of the head may be subjected to pressure from the hip capsule and abductors, resulting in lateral flattening. The combination of medial and lateral flattening can produce a triangular or dunce's cap appearance to the head as seen on an anteroposterior radiograph. Finally, direct force on the head from the reflected head of rectus femoris can produce a groove on the superior part of the head. The end result of the femoral displacement is often loss of articular cartilage and early osteoarthritis. Other directions of dislocation include anterior (8 %) and posterior (1 %), and are sometimes only visible on three-dimensional computed tomographic scans [5]. Increased anteversion and/or valgus and muscle contractures may

J. E. Robb (✉)
School of Medicine, University of St Andrews, St Andrews,
KY16 9TF Scotland, UK
e-mail: j.e.robbs@btinternet.com

G. Hägglund
Department of Orthopaedics, University Hospital,
SE-221 85 Lund, Sweden

also contribute to the femoral deformity [6]. Although the direction of spastic hip subluxation is generally agreed to be posterolateral, a recent radiographic study has suggested that the location of the acetabular deficiency is global and more apparent anteriorly [7].

Hip surveillance

Preventing hip dislocation in a population of children with CP requires a surveillance programme, including routines for preventive treatment of children with hips at risk [8]. Clinical examination of hip displacement in CP is unreliable, and hip range of motion is also a poor indicator of the risk of displacement, as contractures are common and restrict motion. Thus hip displacement can occur in a hip with a normal range of motion [9] and the diagnosis is made from radiographs. Lateral displacement of the femoral head is common without acetabular dysplasia, and acetabular dysplasia develops at a later stage than femoral head lateralization [10]. An early radiographic indicator is a notch appearing on the superolateral aspect of the femoral head, which is thought to be due to moulding in response to the tendency of the femur to migrate proximally and laterally [11]. Measurement of the lateral displacement is sufficient for hip screening in CP [10, 12, 13]. The Reimers' migration percentage [14] (MP) (Fig. 1) is a valid and reliable measure [13–15], and probably the most commonly used. Most reports define a hip with an MP of >30–33 % as displaced and a hip with an MP of >90–100 % as dislocated [12, 14, 16]. Some hips with an MP of <40 % do not progress to dislocation, while almost all hips with an MP of >40 % need preventive surgery to prevent dislocation [10].

The most common age of displacement with an MP of >33 % or >40 % is 3–4 years (Fig. 2), and some children already show displacement at 2 years of age [9]. For an effective surveillance programme, children with CP should be identified early, and those at risk examined radiographically as early as possible.

Hip displacement is related to CP subtype [9]. Children with spastic hemiplegia or ataxic CP have a low risk, and children with spastic tetraplegia or quadriplegia have a high risk. There are, however, several problems with using CP subtype as an indicator of hips at risk in a surveillance programme. There are several different classification systems, and the descriptors for the motor types of CP have not been universally agreed upon; new definitions are under development, and the definitions may be difficult to apply in a reliable manner [17]. It is sometimes impossible to determine the CP subtype before the age of 4 years.

Hip displacement is directly related to the level of gross motor function measured with the gross motor function




Fig. 1 Measurement of the migration percentage (MP). $MP = a/b \times 100$




Fig. 2 Number of children versus age (years) at first registration of an MP above 33 % (yellow) and 40 % (red) in the total population of children with CP




Fig. 3 Proportion of children (%) with an MP of <33 % (green), 33–39 % (yellow) and ≥40 % (red) in relation to GMFCS level in the total population of children with CP

classification scale (GMFCS) [18]. Similar results have been shown from the Victorian Cerebral Palsy Register [19], Sweden [9] and Norway [20] (Fig. 3). The GMFCS

has proven to be a valid and reliable tool [18] and has been reported to remain relatively stable over time [21, 22]. According to the designers of the GMFCS, most children will remain at the same level from the age of two [23]. By using GMFCS instead of CP subtype as an indicator of hips at risk in a surveillance programme, all of the problems described for subtypes seem to be avoided.

In southern Sweden, a follow-up programme for CP was started in 1994 with the specific aim of identifying patients at risk of hip dislocation and preventing this complication. During the subsequent 10 years, the hip dislocation rate dropped from 8 to 0.5 % [16]. Comparable results were reported from an Australian programme introduced in 1997 [12]. Both programmes are based on the child's age and GMFCS level (Fig. 4). The Australian programme contains more frequent radiographic examinations, but in the Swedish programme the children have at least annual clinical examinations by their physiotherapist.

In the Swedish programme, children at GMFCS I are not examined radiographically unless deterioration occurs in the clinical examinations. Children at GMFCS II have radiographic examinations at 2 and 6 years of age, and children at GMFCS III–V have annual examinations from the first suspicion of the diagnosis of CP until 8 years of age. After 8 years of age, the time interval between examinations is assessed individually based on the results of the previous clinical and radiological examinations. In the CPUP programme, all children have clinical examinations twice a year until 6 years of age and once a year thereafter [24].

In the Australian programme, all children have a radiographic examination at 12–24 months of age. Children at GMFCS I are reviewed at 3 and 5 years of age. Children at GMFCS II are examined radiographically once a year until the MP is stable, and all children are examined radiographically at 5 and 8 years of age. Children at

GMFCS III–IV are examined radiographically twice a year until the MP is stable, then once a year until 7 years of age. If the MP is <30 %, radiographic examination is discounted until pre-puberty. The children are examined once a year from pre-puberty until skeletal maturity. Children at GMFCS V have radiographic examinations twice a year until 7 years of age, and then annual examinations until skeletal maturity [25].

Management of hip displacement

The common indications for management of a displaced hip are instability that cannot be controlled by the patient, and restriction of motion and pain, which may already be present in mild subluxations [26]. Functionally, this may manifest as difficulties with transfers, perineal access for hygiene purposes, in obtaining stable seating, and problems with pelvic and spinal alignment. There is controversy, however, as to whether a dislocated hip is painful, but Cooperman et al. [27] suggested that 50 % of a small group of institutionalized patients with CP complained of pain in the dislocated hip. Noonan et al. [28] found that the incidence of hip pain was low and not associated with hip displacement or osteoarthritis in their study of 77 adults with CP. They recommended that surgical treatment of the hip in severely affected patients should be based on the presence of pain or contractures and not on radiographic signs of hip displacement or osteoarthritis.

Non-operative prevention consists of positioning the child to prevent prolonged adduction–flexion at the hip. Optimal lying and sitting positioning during long periods [29] and standing treatment with hips in abduction [30] seem to reduce hip displacement. Bracing is not thought to be effective in spastic hip displacement. Graham et al. [31] studied the combined effect of intramuscular injection of botulinum toxin A and abduction hip bracing in the management of spastic hip displacement in a randomized trial, and found that progressive hip displacement continued to occur in the treatment group, so they did not recommend this treatment. Spasticity-reducing treatment with intrathecal baclofen, selective dorsal rhizotomy or botulinum toxin reduces the adductor spasticity but has not, so far, been proven to reduce the risk of hip displacement.

Operative management to prevent dislocation consists of soft tissue release, varus osteotomy of the proximal femur and pelvic osteotomy. There are still no studies that have shown at what age or at what degree of displacement which of these operations is preferable. The muscle imbalance over the hip joint is present postoperatively and, irrespective of the operative method used, there is a risk of re-displacement during the remaining postoperative growth period.




Fig. 4 Schematic presentation of the radiographic examinations in the Australian and Swedish surveillance programmes. *Green* examination yearly, *red* examination twice a year, *yellow* no examination, *grey* individually based examinations

Soft-tissue release consists of myototomy of the adductor longus, brevis and often gracilis. The operation should also include iliopsoas tenotomy. It has been questioned whether the iliopsoas tendon should be divided at its insertion at the lesser trochanter or by an intramuscular lengthening at the pelvic brim to prevent flexor weakness. Bialic et al. [32], in a comparative study, found more robust results and no adverse effects related to flexor weakness after division at the lesser trochanter. Additionally, division of the psoas tendon at the lesser trochanter in children who do not walk is unlikely to produce an adverse functional effect on hip flexor function. Several studies have shown that children undergoing unilateral soft-tissue release have a high risk of displacement of the contralateral hip [14, 33, 34], and these authors conclude that soft-tissue release should be carried out bilaterally. Adductor longus intramuscular tenotomy, gracilis myotomy and psoas tenotomy is only likely to be effective in the longer term if hip abduction can be maintained. A review of the effects of surgical adductor releases for hip subluxation in cerebral palsy (CP) [35] concluded that there was weak evidence to support the efficacy of the procedure. Interestingly, a very recent report [36] found that adductor surgery was only successful in 27 % of children at GMFCS level IV and 14 % of children at GMFCS level V. The criteria for “success” in this study [36] was defined as the absence of subsequent surgical procedures during the study period and a migration percentage of <50 % in both hips at the time of follow-up.

Progressive uncovering of the femoral head requires the addition of varus derotation osteotomy, provided the acetabulum is not also deformed. A varus osteotomy of the femur has the same effect as passive hip abduction but has the advantage that its effect is independent of passive movement. The femoral deformity is either excessive anteversion or valgus or both. Robin et al. [37] have shown that the neck–shaft angle and anteversion increases with GMFCS level as a result of delayed walking, limited walking or inability to walk. It should be remembered that coxa valga is often a radiographic artefact due to femoral torsion, and a corrected view of the proximal femur should be obtained. An anteroposterior radiograph with the femur in internal rotation to neutralize anteversion is a prerequisite to show the true femoral pathology and the neck–shaft angle. Significant re-operation rates have been reported after early soft tissue or proximal femoral procedures when performed at an average age of 4 years as acetabular remodelling did not reliably occur postoperatively [38]. Others [39] have reported satisfactory results of femoral osteotomy in patients whose pre-operative MP was >50 %. Varus osteotomy, sometimes combined with derotation to reduce increased anteversion of the proximal femur, is made at the level of the lesser trochanter. A further controversy is whether or not the opposite femur should be

osteotomized, and some authors consider that only the displaced hip should be operated on [36, 38, 39], whereas others recommend bilateral osteotomy in all cases [40].

Where there is a combination of acetabular and femoral head deformity, the aim of surgery is to relocate the hip by reducing the neck–shaft angle to about 120°, anteversion to a physiological value, and to cover the femoral head with a pelvic osteotomy such as a Pemberton [41] or Dega type of periacetabular osteotomy [42]. The Dega type, as originally described [42], extended into the sciatic notch, but many surgeons direct the osteotomy down the posterior column towards the posterior part of the triradiate cartilage [43, 44]. It is also possible to perform a Dega type of osteotomy after closure of the triradiate cartilage in patients with severe CP, as the pelvic bone is often osteoporotic [45]. An open reduction of the hip is not always necessary, but when needed should include resection of the ligamentum teres and division of the transverse acetabular ligament. Sufficient shortening of the femur at the time of femoral osteotomy is most important to lengthen, relatively, the contracted muscles and to avoid excessive pressure on the femoral head; additional soft tissue procedures are not usually necessary as part of this procedure. Anterior hip dislocation can pose a challenge and a Pemberton osteotomy, rather than a Dega type, may give a better prospect of anterior coverage. Even severe deformity of the femoral head is not necessarily a contraindication to reduction, since the risk of subsequent symptomatic arthritis appears to be small.

The evaluation of functional outcomes is challenging in patients with severe CP who have undergone surgery for hip displacement. The literature consists mainly of case series of relatively small numbers of patients with short- to medium-term follow-up, and not always conformity of management or outcome measures between case series. However, it seems that a one-stage hip reduction and pelvic and femoral osteotomy is generally successful [46] and gives superior results to femoral osteotomy alone [39, 47]. In experienced hands, reconstructions for lateral or posterior dislocations have given success rates of 95 % for range of motion and hip reduction and 90 % for pain reduction [48].

The unreconstructable hip

The demand on the hip in patients with severe CP is low and osteoarthritic changes seen radiologically can remain asymptomatic for many years. Hip pain does not seem to correlate well with femoral head deformity. If the hip is symptomatic and unreconstructable, there is a choice of salvage procedures to obtain reasonable abduction and relief of pain. These include proximal femoral resection [49–61], prosthetic replacement of the hip [62–70],

prosthetic interposition arthroplasty [71], valgus femoral osteotomy [72, 73], a combination of valgus osteotomy and proximal femoral excision [55, 57], and hip fusion [66].

The commonest reported procedure is a proximal femoral resection described by Castle and Schneider [49]. The gluteus medius, minimus and vastus lateralis are used as an interposition between the resected femur and acetabulum. About 80 % of patients are comfortable afterwards but insufficient femoral resection and heterotopic bone formation are two important potential complications of the procedure. A proximal femoral excision is not recommended for patients who have a displaced hip but are capable of walking, or have independent or assisted transfer capability, because of the subsequent loss of hip stability. In a recent report, Egermann et al. [50] have reported a reduction in heterotopic bone formation from about 80 to 60 % after a Castle and Schneider [49] procedure by “capping” the proximal femoral stump with the excised femoral head.

Although the Castle and Schneider [49] procedure is the most commonly reported solution for the painful unreconstructable hip, the literature on salvage procedures consists of mainly case series with small numbers and short-term follow-ups, and, again, outcome measures are difficult to evaluate in patients who have severe CP.

Conclusion

Hip dislocation in CP is potentially preventable if children are included from an early age in a surveillance programme that includes repeat radiographic and clinical examinations and preventive treatment for hips that are displacing. The surveillance programme should be based on the child's age, GMFCS level and the degree of lateral displacement as measured by the MP. Non-operative prevention by proper positioning should be considered in all children with CP, although the evidence base for this is weak. Operative prevention should be considered in children with an MP exceeding 33 %. Children will require rehabilitation after any surgical procedure, and this should be factored into any decision about surgery, which would also take into account the wishes of the child and/or their family/carers. Children should also be followed postoperatively until at least skeletal maturity, and if the child has scoliosis associated with a pelvic tilt they should, ideally, be followed further into adulthood.

References

1. Tachdjian MO, Minear WL (1956) Hip dislocation in cerebral palsy. *J Bone Joint Surg Am* 38A:1358–1364
2. Miller F (2005) In: Cerebral palsy. Springer, Berlin, pp 524–525
3. Brown JK, Minns RA (1989) Mechanisms of deformity in children with cerebral palsy. *Sem Orthop* 4:236–255
4. Sanger TD, Kukke SN (2007) Abnormalities of tactile sensory function in children with dystonic and diplegic cerebral palsy. *J Child Neurol* 22:289–293
5. Brunner R, Picard C, Robb JE (1997) Morphology of the acetabulum in hip dislocations due to cerebral palsy. *J Pediatr Orthop B* 6:207–211
6. Brunner R, Baumann JU (1997) Long-term effects of intertrochanteric varus-derotation osteotomy on femur and acetabulum in spastic cerebral palsy. *J Pediatr Orthop* 17:585–591
7. Chang CH, Kuo KN, Wang CJ, Chen YY, Cheng HY, Kao HK (2011) Acetabular deficiency in spastic hip subluxation. *J Pediatr Orthop* 31:648–654
8. Gordon GS, Simkiss DE (2006) A systematic review of the evidence for hip surveillance in children with cerebral palsy. *J Bone Joint Surg Br* 88B:1492–1496
9. Häggglund G, Lauge-Pedersen H, Wagner P (2007) Characteristics of children with hip displacement in cerebral palsy. *BMC Musculoskelet Disord* 8:101
10. Häggglund G, Lauge-Pedersen H, Persson M (2007) Radiographic threshold values for hip screening in cerebral palsy. *J Child Orthop* 1:43–47
11. Howard CB, Williams LA (1984) A new radiological sign in the hips of cerebral palsy patients. *Clin Radiol* 35:317–319
12. Dobson F, Boyd RN, Parrott J, Nattrass GR, Graham HK (2002) Hip surveillance in children with cerebral palsy. *J Bone Joint Surg Br* 84B:720–726
13. Parrott J, Boyd RN, Dobson F, Lancaster A, Love S, Oates J, Wolfe R, Nattrass GR, Graham HK (2002) Hip displacement in spastic cerebral palsy: repeatability of radiologic measurement. *J Pediatr Orthop* 22:660–667
14. Reimers J (1980) The stability of the hip in children: a radiological study of the results of muscle surgery in cerebral palsy. *Acta Orthop Scand Suppl* 184:1–100
15. Faraj S, Atherton WG, Stott NS (2004) Inter- and intra-measurer error in the measurement of Reimer's hip migration percentage. *J Bone Joint Surg Br* 86B:434–437
16. Häggglund G, Andersson S, Dümpe H, Lauge Pedersen H, Nordmark E, Westbom L (2005) Prevention of dislocation of the hip in children with cerebral palsy. First 10 years experience of a population based prevention program. *J Bone Joint Surg Br* 87B:95–101
17. Stanley F, Blair E, Alberman E (2000) Cerebral palsies: epidemiology and causal pathways. Mac Keith, London, pp 14–21
18. Palisano R, Rosenbaum P, Walter S, Russell D, Wood E, Galuppi B (1997) Development and reliability of a system to classify gross motor function in children with cerebral palsy. *Dev Med Child Neurol* 39:214–223
19. Soo B, Howard JJ, Boyd R, Reid SM, Lanigan A, Wolfe R, Reddiough D, Graham K (2006) Hip displacement in cerebral palsy. *J Bone Joint Surg Am* 88A:121–129
20. Elkamil AI, Andersen GL, Häggglund G, Lamvik T, Skranes J, Vik T (2011) Prevalence of hip dislocation in children with cerebral palsy in regions with and without a surveillance programme. A cross sectional study in Sweden and Norway. *BMC Musculoskelet Disord* 12:284
21. Wood E, Rosenbaum P (2000) The gross motor function classification system for cerebral palsy: a study of reliability and stability over time. *Dev Med Child Neurol* 42:292–296
22. Palisano R, Cameron D, Rosenbaum P, Walter SD, Russel D (2006) Stability of the gross motor function classification system. *Dev Med Child Neurol* 48:424–428
23. Rosenbaum P, Walter SD, Hanna SE, Palisano RJ, Russel DJ, Parminder R, Wood E, Bartlett DJ, Galuppi B (2002) Prognosis

- for gross motor function in children with cerebral palsy: creation of motor development curves. *JAMA* 288:1357–1363
24. Hedberg A et al. (2013) CPUP website. www.cpunet.se
 25. Australasian Academy of Cerebral Palsy and Developmental Medicine (2013) Website. www.ausacpdm.org.au
 26. Brunner R, Baumann JU (1994) Clinical benefit of reconstruction of dislocated and subluxated hip joints in patients with spastic cerebral palsy. *J Pediatr Orthop* 14:290–294
 27. Cooperman DR, Bartucci E, Dierrick E, Millar EA (1987) Hip dislocation in spastic cerebral palsy: long-term consequences. *J Pediatr Orthop* 7:268–276
 28. Noonan KJ, Jones J, Pierson J, Honkamp NJ, Leverson G (2004) Hip function in adults with severe cerebral palsy. *J Bone Joint Surg Am* 86A:2607–2613
 29. Poutney TE, Mandy A, Green E, Gard PR (2009) Hip subluxation and dislocation in cerebral palsy: a prospective study on the effectiveness of postural management programmes. *Physiother Res Int* 14:116–127
 30. Martinsson C, Himmelmann K (2011) Effect of weight-bearing in abduction and extension on hip stability in children with cerebral palsy. *Pediatr Phys Ther* 23:150–157
 31. Graham HK, Boyd RN, Carlin JB, Dobson F, Lowe K, Nattrass G, Thomason P, Wolfe R, Reddiough D (2008) Does botulinum toxin A combined with bracing prevent hip displacement in children with cerebral palsy and ‘hips at risk’? A randomized, controlled trial. *J Bone Joint Surg Am* 90A:23–33
 32. Bialik GM, Pierce R, Dorociak R, Lee TS, Aiona MD, Susman MD (2009) Iliopsoas tenotomy at the lesser trochanter versus at the pelvic brim in ambulatory children with cerebral palsy. *J Pediatr Orthop* 29:252–255
 33. Samilson RL, Carson JJ, James P, Raney FL Jr (1967) Results and complications of adductor tenotomy and obturator neurectomy in cerebral palsy. *Clin Orthop Relat Res* 54:61–73
 34. Silver RL, Rang M, Chan J, de la Garza J (1985) Adductor release in non-ambulant children with cerebral palsy. *J Pediatr Orthop* 5:672–677
 35. Stott NS, Piedrahita L (2004) Effects of surgical adductor releases for hip subluxation in cerebral palsy: an AACPD evidence report. *Dev Med Child Neurol* 46:628–645
 36. Shore BJ, Yu X, Desai S, Selber P, Wolfe R, Graham HK (2012) Adductor surgery to prevent hip displacement in children with cerebral palsy: the predictive role of the gross motor function classification system. *J Bone Joint Surg Am* 94A:326–334
 37. Robin J, Graham HK, Selber P, Dobson F, Smith K, Baker R (2008) Proximal femoral geometry in cerebral palsy. A population based cross sectional study. *J Bone Joint Surg Br* 90B:1372–1379
 38. Schmale GA, Eilert RE, Chang F, Seidel K (2006) High reoperation rates after early treatment of the subluxating hip in children with spastic cerebral palsy. *J Pediatr Orthop* 26:617–623
 39. Huh K, Rethlefsen SA, Wren TA, Kay RM (2011) Surgical management of hip subluxation and dislocation in children with cerebral palsy: isolated VDRO or combined surgery? *J Pediatr Orthop* 8:858–863
 40. Owers KL et al (2001) Bilateral hip surgery in severe cerebral palsy a preliminary review. *J Bone Joint Surg Br* 83B:1161–1167
 41. Pemberton PA (1965) Pericapsular osteotomy of the ilium for treatment of congenital subluxation and dislocation of the hip. *J Bone Joint Surg Am* 47A:65–86
 42. Dega W (1974) Transiliac osteotomy in the treatment of congenital hip dysplasia. *Chir Narzadow Ruchu Ortop Pol* 39:601–613
 43. Mubarak SJ, Valencia FG, Wenger DR (1992) One-stage correction of the spastic dislocated hip. Use of pericapsular acetabuloplasty to improve coverage. *J Bone Joint Surg Am* 74A:1347–1357
 44. Miller F, Girardi H, Lipton G, Ponzio R, Klaumann M, Dabney KW (1997) Reconstruction of the dysplastic spastic hip with periliac pelvic and femoral osteotomy followed by immediate mobilization. *J Pediatr Orthop* 17:592–602
 45. Robb JE, Brunner R (2006) A Dega-type osteotomy after closure of the triradiate cartilage in non-walking patients with severe cerebral palsy. *J Bone Joint Surg Br* 88B:933–937
 46. Sankar WN, Spiegel DA, Gregg JR, Sennett BJ (2006) Long-term follow up after one stage reconstruction of dislocated hips in patients with cerebral palsy. *J Pediatr Orthop* 26:1–7
 47. Al-Ghadir M, Masquijo JJ, Guerra LA, Willis B (2009) Combined femoral and pelvic osteotomies versus femoral osteotomy alone in the treatment of hip dysplasia in children with cerebral palsy. *J Pediatr Orthop* 29:779–783
 48. Miller F (2005) In: *Cerebral palsy*. Springer, Berlin, pp 550–551
 49. Castle ME, Schneider C (1978) Proximal femoral resection-interposition arthroplasty. *J Bone Joint Surg Am* 60A:1051–1054
 50. Egermann M, Döderlein L, Schlager E, Müller S, Braatz F (2009) Autologous capping during resection arthroplasty of the hip in patients with cerebral palsy. *J Bone Joint Surg Br* 91B:1007–1012
 51. Knaus A, Terjesen T (2009) Proximal femoral resection arthroplasty for patients with cerebral palsy and dislocated hips: 20 patients followed for 1–6 years. *Acta Orthop* 80:32–36
 52. Muthusamy K, Chu HY, Friesen RM, Chou PC, Eilert RE, Chang FM (2008) Femoral head resection as a salvage procedure for the severely dysplastic hip in non-ambulatory children with cerebral palsy. *J Pediatr Orthop* 28:884–889
 53. Lampropoulos M, Puigdevall MH, Zapozko D, Malvárez HR (2008) Proximal femoral resection and articulated hip distraction with an external fixator for the treatment of painful spastic hip dislocations in paediatric patients with spastic quadriplegia. *J Pediatr Orthop B* 17:27–31
 54. Abu-Rajab RB, Bennet GC (2007) Proximal femoral resection-interposition arthroplasty in cerebral palsy. *J Pediatr Orthop B* 16:181–184
 55. Leet AI, Chhor K, Launay F, Kier-York J, Sponseller PD (2005) Femoral head resection for painful hip subluxation in cerebral palsy: is valgus osteotomy in conjunction with femoral head resection preferable to proximal femoral head resection and traction? *J Pediatr Orthop* 25:70–73
 56. Widmann RF, Do TT, Doyle SM, Burke SW, Root L (1999) Resection arthroplasty of the hip for patients with cerebral palsy: an outcome study. *J Pediatr Orthop* 19:805–810
 57. McHale KA, Bagg M, Nason SS (1990) Treatment of the chronically dislocated hip in adolescents with cerebral palsy with femoral head resection and subtrochanteric valgus osteotomy. *J Pediatr Orthop* 10:504–509
 58. McCarthy RE, Simon S, Douglas B, Zawacki R, Reese N (1988) Proximal femoral resection to allow adults who have severe cerebral palsy to sit. *J Bone Joint Surg Am* 70A:1011–1016
 59. Baxter MP, D’Astous JL (1986) Proximal femoral resection-interposition arthroplasty: salvage hip surgery for the severely disabled child with cerebral palsy. *J Pediatr Orthop* 6:681–685
 60. Koffman M (1981) Proximal femoral resection or total hip replacement in severely disabled cerebral-spastic patients. *Orthop Clin North Am* 12:91–100
 61. Perlmutter MN, Synder M, Miller F, Bisbal R (1993) Proximal femoral resection for older children with spastic hip disease. *Dev Med Child Neurol* 35:525–531
 62. Queally JM, Abdulkarim A, Mulhall KJ (2009) Total hip replacement in patients with neurological conditions. *J Bone Joint Surg Br* 91B:1267–1273
 63. Schroeder K, Hauck C, Wiedenhöfer B, Braatz F, Aldinger PR (2010) Long-term results of hip arthroplasty in ambulatory patients with cerebral palsy. *Int Orthop* 34:335–339

64. Blake SM, Kitson J, Howell JR, Gie GA, Cox PJ (2006) Constrained total hip arthroplasty in a paediatric patient with cerebral palsy and painful dislocation of the hip. A case report. *J Bone Joint Surg Br* 88B:655–657
65. Weber M, Cabanela ME (1999) Total hip arthroplasty in patients with cerebral palsy. *Orthopedics* 22:425–427
66. Root L, Goss JR, Mendes J (1986) The treatment of the painful hip in cerebral palsy by total hip replacement or hip arthrodesis. *J Bone Joint Surg Am* 68A:590–598
67. Skoff HD, Keggi K (1986) Total hip replacement in the neuro-muscularly impaired. *Orthop Rev* 15:154–159
68. Root L (1982) Total hip replacement in young people with neurological disease. *Dev Med Child Neurol* 24:186–188
69. Buly RL, Huo M, Root L, Binzer T, Wilson PD Jr (1993) Total hip arthroplasty in cerebral palsy. Long-term follow-up results. *Clin Orthop Relat Res* 296:148–153
70. Prosser GH, Shears E, O'Hara JN (2012) Hip resurfacing with femoral osteotomy for painful subluxed or dislocated hips in patients with cerebral palsy. *J Bone Joint Surg Br* 94B:483–487
71. Gabos PG, Miller F, Galban MA, Gupta GG, Dabney K (1999) Prosthetic interposition arthroplasty for the palliative treatment of end-stage spastic hip disease in nonambulatory patients with cerebral palsy. *J Pediatr Orthop* 19:796–804
72. Schejbalova A, Havlas V, Trc T (2009) Irreducible dislocation of the hip in cerebral palsy patients treated by Schanz proximal femoral valgus osteotomy. *Int Orthop* 33:1713–1717
73. Hogan KA, Blake M, Gross RH (2006) Subtrochanteric valgus osteotomy for chronically dislocated, painful spastic hips. *J Bone Joint Surg Am* 88A:2624–2631